

Eymox



Indication	Bacterial conjunctivitis, tarsadenitis, keratitis, corneal ulcer, and also as an aseptic measure in ophthalmic surgeries
Effective Bacillus Strain	<i>Staphylococcus</i> , <i>Streptococcus</i> , <i>Streptococcus pneumonia</i> , <i>Micrococcus</i> , <i>Enterococci</i> , <i>Corynebacterium spp.</i> , <i>Citrobacter spp.</i> , <i>Klebsiella spp.</i> , <i>Enterobacter spp.</i> , <i>Serratia spp.</i> , <i>Proteus spp.</i> , <i>Morganella morganii</i> , <i>influenza bacteria</i> , <i>Pseudomonas aeruginosa</i> , <i>Burkholderia cephalcy</i> , <i>Stenotropomonas (Santomonas) Maltophilia</i> , <i>Acinetobacter spp.</i> , <i>Propionibacterium acnes</i>
Active Ingredient	Moxifloxacin Hydrochloride 5.45 mg/mL
Description	Light greenish yellow ophthalmic solution
Dosage & Administration	<p>Bacterial conjunctivitis, tarsadenitis, keratitis, corneal ulcer: Instill 1 drop into the affected eye(s) 3 times daily</p> <p>Pre-operation: Instill 1 drop into the affected eye(s) 5 times daily</p> <p>Post-operation: Instill 1 drop into the affected eye(s) 3 times daily</p>
Package	5 mL/bottle